

Supersymmetric Electroweak Corrections to the Chargino Decay into Neutralino and W Boson *

Zhang Ren-You², Ma Wen-Gan^{1,2}, Wan Lang-Hui²

¹ CCAST (World Laboratory), P.O.Box 8730, Beijing 100080, P.R.China

² Department of Modern Physics, University of Science and Technology
of China (USTC), Hefei, Anhui 230027, P.R.China

Abstract

We investigated the electroweak corrections from one-loop diagrams involving the third generation (s)quarks to the decay width of process $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$ in the framework of the Minimal Supersymmetric Standard Model. Our calculation shows that these corrections are not very sensitive to the mass of the lightest neutralino $\tilde{\chi}_1^0$ and the sbottom mixing angle θ_b , but depend strongly on the top squark mixing angle θ_t and $\tan\beta$. With our chosen parameters, we find that these radiative corrections can exceed 10%, therefore they should be taken into account for the precise experimental measurement at future colliders.

PACS: 11.30.Pb, 12.15.Lk, 12.60.Jv, 14.80.Ly

*Supported by National Natural Science Foundation of China.

I Introduction

Over the past years, many experimental works were focused on the finding new physics beyond the standard model (SM) [1]. At the same time, some new theoretical models have been developed in order to describe the new physics, such as the minimal supersymmetric model (MSSM) [2] [3] [4], large extra dimensions model [5], etc. Among these models, the MSSM is arguably the most promising candidate for extension beyond the SM, in which all the SM particles have their supersymmetric partners. The existence of these supersymmetric particles is one of the characteristics of this theory. Therefore, searching for these supersymmetric particles at colliders is an important task to verify the MSSM. In the MSSM there exist two charginos $\tilde{\chi}_i^+$ ($i = 1, 2$) and four neutralinos $\tilde{\chi}_i^0$ ($i = 1, \dots, 4$). They are just the mixtures of gauginos and higgsinos, the supersymmetric partners of gauge bosons and Higgs bosons [2] [3]. The lightest neutralinos $\tilde{\chi}_1^0, \tilde{\chi}_2^0$ and chargino $\tilde{\chi}_1^+$ ($m_{\tilde{\chi}_1^+} < m_{\tilde{\chi}_2^+}$) are expected to be the lightest supersymmetric particles. For R-parity conservation, the lightest supersymmetric particle (LSP) $\tilde{\chi}_1^0$ is stable and invisible. Thus, the states $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^+$ might be the first supersymmetric particles to be discovered. The search for $\tilde{\chi}_1^+$ and $\tilde{\chi}_2^0$ is the main task at present and future colliders. Many experiments have been done at present colliders, such as LEP, Tevatron, etc., in searching for the supersymmetric particles [6] [7] [8], especially for charginos and neutralinos [9] [10]. At the same time, the phenomenological calculations of chargino and neutralino productions via $\gamma\gamma$, $e^-\gamma$, e^+e^- , e^-e^- , γp , pp collisions, etc., have also been performed. They can be found in Ref.[11, 12, 13, 14, 15, 16, 17].

The decays of charginos and neutralinos in the MSSM have been widely discussed

[18] [19]. If the mass splitting between $\tilde{\chi}_1^0$ (LSP) and $\tilde{\chi}_1^+$ or $\tilde{\chi}_2^0$ is not large enough, the three-body decay modes of $\tilde{\chi}_1^+$ and $\tilde{\chi}_2^0$ into $\tilde{\chi}_1^0$ and two fermions through vector boson, Higgs boson and sfermion exchanges are the dominant decay modes, which take place mainly in the minimal supergravity (MSUGRA) scenario. But in the models without gaugino mass unification at the GUT scale, the mass splitting between them can exceed the masses of gauge bosons (W, Z) and Higgs bosons in some parameter space. Therefore, the decay modes $\tilde{\chi}_1^+(\tilde{\chi}_2^0) \rightarrow W^+(Z^0) + \tilde{\chi}_1^0$, $\tilde{\chi}_1^+(\tilde{\chi}_2^0) \rightarrow H^+(h^0, H^0, A^0) + \tilde{\chi}_1^0$ are kinematically accessible. In this case, the above two-body decay channels will be the most important decay modes of $\tilde{\chi}_1^+$ and $\tilde{\chi}_2^0$, since the chargino and neutralino have enough phase space to decay firstly via these channels. If $\tilde{\chi}_1^+$ and $\tilde{\chi}_2^0$ are rather heavy, some new decay channels, such as decay into $f\tilde{f}$ pair, should be under consideration when they are kinematically allowed [18]. In this paper, we will only consider the decay of the lightest chargino into the lightest neutralino and W boson. As mentioned above, in the case of $m_{\tilde{\chi}_1^+} > m_W + m_{\tilde{\chi}_1^0}$, this decay channel can open up and will be the dominant decay channel of $\tilde{\chi}_1^+$. The W boson's leptonic decay may be used as a spectacular trigger. For the comparison of the theoretical prediction with the precise measurement at the future colliders, the more accurate calculation of the partial decay width of this decay mode is necessary. For this purpose, the radiative corrections at one-loop level should be included. Since there is no one-loop QCD correction to this process, the electroweak corrections from one-loop quark-squark diagrams are the most important. In this paper, only the third generation quarks and squarks are considered because of the Yukawa characteristic factor m_q/m_W in the chargino/neutralino-quark-squark vertices (m_q is the mass of quark q)[20]. In our

calculation we adopted the t'Hooft gauge. The relevant diagrams of the tree-level, counter-term, one-loop vertex corrections to this decay and the self-energies of W-boson, charginos and neutralinos, are shown in Fig.1(a), (b), (c), (d), (e) and (f) ($i = 1, j = 1$), respectively.

The paper is organized as follows: In section 2, we review briefly the chargino, neutralino and squark sectors of the MSSM and give the tree-level results (amplitude and decay width). In section 3, we give some analytical results of the electroweak corrections to the $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$ partial decay width. The numerical results and our conclusions are presented in section 4.

II Notations and tree-level results

The tree-level Lagrangian for the chargino and neutralino masses reads

$$\mathcal{L}_m^{\tilde{\chi}} = -\frac{1}{2} \begin{pmatrix} \psi^+ \\ \psi^- \end{pmatrix}^T \begin{pmatrix} 0 & X^T \\ X & 0 \end{pmatrix} \begin{pmatrix} \psi^+ \\ \psi^- \end{pmatrix} - \frac{1}{2} \psi^{0T} Y \psi^0 + h.c. \quad (2.1)$$

where the tree-level mass matrices X, Y are [20]

$$\begin{aligned} X &= \begin{pmatrix} M & \sqrt{2}m_W \sin \beta \\ \sqrt{2}m_W \cos \beta & \mu \end{pmatrix} \\ Y &= \begin{pmatrix} M' & 0 & -m_Z s_W \cos \beta & m_Z s_W \sin \beta \\ 0 & M & m_Z c_W \cos \beta & -m_Z c_W \sin \beta \\ -m_Z s_W \cos \beta & m_Z c_W \cos \beta & 0 & -\mu \\ m_Z s_W \sin \beta & -m_Z c_W \sin \beta & -\mu & 0 \end{pmatrix} \end{aligned} \quad (2.2)$$

and $\psi^{+T} = (-i\lambda^+, \psi_{H_2}^+)$, $\psi^{-T} = (-i\lambda^-, \psi_{H_1}^-)$, $\psi^{0T} = (-i\lambda', -i\lambda^3, \psi_{H_1}^0, \psi_{H_2}^0)$, where $s_W \equiv \sin \theta_W$, $c_W \equiv \cos \theta_W$, M and M' are the $SU(2)$ and $U(1)_Y$ soft-SUSY-breaking parameters, respectively. We define

$$\chi^+ = V\psi^+,$$

$$\chi^- = U\psi^-,$$

$$\chi^0 = N\psi^0, \quad (2.3)$$

where U , V and N are unitary matrices chosen such that:

$$\begin{aligned} U^* X V^{-1} &= M_D = \text{diag}(m_{\tilde{\chi}_1^+}, m_{\tilde{\chi}_2^+}), \quad (0 < m_{\tilde{\chi}_1^+} < m_{\tilde{\chi}_2^+}) \\ N^* Y N^{-1} &= M_D^0 = \text{diag}(m_{\tilde{\chi}_1^0}, m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_3^0}, m_{\tilde{\chi}_4^0}), \quad (0 < m_{\tilde{\chi}_1^0} < m_{\tilde{\chi}_2^0} < m_{\tilde{\chi}_3^0} < m_{\tilde{\chi}_4^0}). \end{aligned} \quad (2.4)$$

The tree-level Lagrangian for the stop and sbottom masses is

$$\mathcal{L}_m^{\tilde{q}} = - \begin{pmatrix} \tilde{t}_L \\ \tilde{t}_R \end{pmatrix}^\dagger \mathcal{M}_t^2 \begin{pmatrix} \tilde{t}_L \\ \tilde{t}_R \end{pmatrix} - \begin{pmatrix} \tilde{b}_L \\ \tilde{b}_R \end{pmatrix}^\dagger \mathcal{M}_b^2 \begin{pmatrix} \tilde{b}_L \\ \tilde{b}_R \end{pmatrix} \quad (2.5)$$

in which the tree-level stop and sbottom squared-mass matrices are:

$$\begin{aligned} \mathcal{M}_t^2 &= \begin{pmatrix} M_{\tilde{Q}}^2 + m_t^2 + m_Z^2 \cos 2\beta \left(\frac{1}{2} - \frac{2}{3}s_W^2\right) & m_t (A_t - \mu \cot \beta) \\ m_t (A_t - \mu \cot \beta) & M_{\tilde{U}}^2 + m_t^2 + \frac{2}{3}m_Z^2 \cos 2\beta s_W^2 \end{pmatrix} \\ \mathcal{M}_b^2 &= \begin{pmatrix} M_{\tilde{Q}}^2 + m_b^2 - m_Z^2 \cos 2\beta \left(\frac{1}{2} - \frac{1}{3}s_W^2\right) & m_b (A_b - \mu \tan \beta) \\ m_b (A_b - \mu \tan \beta) & M_{\tilde{D}}^2 + m_b^2 - \frac{1}{3}m_Z^2 \cos 2\beta s_W^2 \end{pmatrix} \end{aligned} \quad (2.6)$$

where $M_{\tilde{Q}}$, $M_{\tilde{U}}$ and $M_{\tilde{D}}$ are the soft-SUSY-breaking masses. For the third generation SU(2) squark doublet, we have $\tilde{Q} = (\tilde{t}_L, \tilde{b}_L)$, and the singlets $\tilde{U} = \tilde{t}_R$ and $\tilde{D} = \tilde{b}_R$, respectively. $A_{t,b}$ are the corresponding soft-SUSY-breaking trilinear coupling parameters.

To diagonalize the mass matrices $\mathcal{M}_t^2$ and $\mathcal{M}_b^2$, we should introduce two unitary matrices $\mathcal{R}^{\tilde{q}}$: $\mathcal{R}^{\tilde{q}} \mathcal{M}_{\tilde{q}}^2 \mathcal{R}^{\tilde{q}+} = \text{diag}(m_{q_1}^2, m_{q_2}^2)$, ($q = t, b$). In consideration of the reality of the mass matrices, we may choose matrices $\mathcal{R}^{\tilde{q}}$ to be

$$\mathcal{R}^{\tilde{q}} = \begin{pmatrix} \cos \theta_q & \sin \theta_q \\ -\sin \theta_q & \cos \theta_q \end{pmatrix} \quad (q = t, b), \quad (2.7)$$

where $-\pi/2 \leq \theta_{\tilde{q}} \leq \pi/2$. The mass-eigenstates $\tilde{q}_1, \tilde{q}_2$ are related to the current eigenstates

$\tilde{q}_L, \tilde{q}_R$ by the transformation:

$$\begin{pmatrix} \tilde{q}_1 \\ \tilde{q}_2 \end{pmatrix} = \mathcal{R}^{\tilde{q}} \begin{pmatrix} \tilde{q}_L \\ \tilde{q}_R \end{pmatrix}. \quad (2.8)$$

The tree-level Lagrangian for the chargino-neutralino-W boson interactions is [3]

$$\mathcal{L}_{\tilde{\chi}^+ \tilde{\chi}^0 W} = g \tilde{\chi}_i^0 \gamma^\mu [O_{ij}^L P_L + O_{ij}^R P_R] \tilde{\chi}_j^+ W_\mu^- + h.c. \quad (2.9)$$

where $P_{L,R} = \frac{1}{2}(1 \mp \gamma_5)$, and

$$\begin{aligned} O_{ij}^L &= -\frac{1}{\sqrt{2}} N_{i4} V_{j2}^* + N_{i2} V_{j1}^* \\ O_{ij}^R &= \frac{1}{\sqrt{2}} N_{i3}^* U_{j2} + N_{i2}^* U_{j1} \end{aligned} \quad (2.10)$$

From this Lagrangian, we can easily obtain the tree-level amplitude and partial width of the decay $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$:

$$\begin{aligned} \Gamma_{tree}(\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0) &= \frac{\lambda}{16\pi m_{\tilde{\chi}_1^+}^3} |\mathcal{M}_{tree}|^2 \\ &= \frac{\lambda g^2}{16\pi m_{\tilde{\chi}_1^+}^3} \left[\rho (|O_{11}^L|^2 + |O_{11}^R|^2) - 12m_{\tilde{\chi}_1^+} m_{\tilde{\chi}_1^0} \text{Re}(O_{11}^{L*} O_{11}^R) \right] \end{aligned} \quad (2.11)$$

where

$$\begin{aligned} \lambda &= \left[\left(m_{\tilde{\chi}_1^+}^2 - (m_{\tilde{\chi}_1^0} + m_W)^2 \right) \left(m_{\tilde{\chi}_1^+}^2 - (m_{\tilde{\chi}_1^0} - m_W)^2 \right) \right]^{1/2} \\ \rho &= m_{\tilde{\chi}_1^+}^2 + m_{\tilde{\chi}_1^0}^2 - 2m_W^2 + \frac{(m_{\tilde{\chi}_1^+}^2 - m_{\tilde{\chi}_1^0}^2)^2}{m_W^2} \end{aligned} \quad (2.12)$$

III Renormalization and radiative corrections

The supersymmetric radiative corrections to chargino and neutralino productions have been considered widely [21] [22]. For convenience, the corrections are expressed by using

form factors. All the one-loop level contributions to these form factors can be classified in terms of prototypes distinguished by the number of particles inside the loops and their spin [22]. The calculation of the supersymmetric corrections to the decay $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$ considered in this paper can be performed analogously. The form factors can be divided into two parts, which are just the contributions of counter-term and vertex correction. In order to obtain the counter-term of vertex $\tilde{\chi}_i^0 \tilde{\chi}_j^+ W^-$ which will be used in the calculation of the corrections, the following renormalization constants are defined

(1) renormalization constants of fields [23]:

$$\begin{aligned} W_\mu &\rightarrow Z_W^{1/2} W_\mu \\ P_L \tilde{\chi}_i^+ &\rightarrow Z_{+,ij}^{L1/2} P_L \tilde{\chi}_j^+, \quad P_R \tilde{\chi}_i^+ \rightarrow Z_{+,ij}^{R1/2} P_R \tilde{\chi}_j^+ \\ P_L \tilde{\chi}_i^0 &\rightarrow Z_{0,ij}^{L1/2} P_L \tilde{\chi}_j^0, \quad P_R \tilde{\chi}_i^0 \rightarrow Z_{0,ij}^{R1/2} P_R \tilde{\chi}_j^0 \end{aligned}$$

(2) renormalization constants of gauge couplings and masses [23]:

$$\begin{aligned} e &\rightarrow Z_e e = (1 + \delta Z_e) e \\ g &\rightarrow g + \delta g = (1 + \delta Z_e - \frac{\delta s_W}{s_W}) g \\ m_W^2 &\rightarrow m_W^2 + \delta m_W^2, \quad m_Z^2 \rightarrow m_Z^2 + \delta m_Z^2 \end{aligned}$$

(3) renormalization of matrices U, V and N [24]:

$$U \rightarrow U + \delta U, \quad V \rightarrow V + \delta V, \quad N \rightarrow N + \delta N, \quad (3.1)$$

where $Z_{0,ij}^R = Z_{0,ij}^{L*}$, because neutralinos $\tilde{\chi}_i^0$ are Majorana particles, $\tilde{\chi}_i^{0c} = \tilde{\chi}_i^0$.

In this paper, we adopt the complete on-mass-shell scheme in doing renormalization.

The relevant renormalization constants in eq.(3.1) are fixed by the on-mass-shell renormalization conditions:

(a) The on-mass-shell renormalization conditions for gauge bosons are [23]

$$\begin{aligned}
\lim_{k^2 \rightarrow m_W^2} \frac{1}{-i(k^2 - m_W^2)} \tilde{R}e i \hat{\Gamma}_{\mu\nu}^{WW}(k) \epsilon^\nu(k) &= \epsilon_\mu(k) \\
\lim_{k^2 \rightarrow m_Z^2} \frac{1}{-i(k^2 - m_Z^2)} \tilde{R}e i \hat{\Gamma}_{\mu\nu}^{ZZ}(k) \epsilon^\nu(k) &= \epsilon_\mu(k) \\
\lim_{k^2 \rightarrow 0} \frac{1}{-i k^2} \tilde{R}e i \hat{\Gamma}_{\mu\nu}^{\gamma\gamma}(k) \epsilon^\nu(k) &= \epsilon_\mu(k) \\
\tilde{R}e i \hat{\Gamma}_{\mu\nu}^{\gamma Z}(k) \epsilon^\nu(k) \big|_{k^2=0, m_Z^2=0} &= 0
\end{aligned} \tag{3.2}$$

where $\epsilon(k)$ are the polarization vectors of the external fields. $\tilde{R}e$ takes only the real part of the loop integrals appearing in the renormalized one-particle-irreducible two-point Green functions $i \hat{\Gamma}_{\mu\nu}$ which can be written as

$$\begin{aligned}
i \hat{\Gamma}_{\mu\nu}^{WW}(k) &= -i g_{\mu\nu} (k^2 - m_W^2) - i \left(g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) \hat{\Sigma}_T^{WW}(k^2) - i \frac{k_\mu k_\nu}{k^2} \hat{\Sigma}_L^{WW}(k^2) \\
i \hat{\Gamma}_{\mu\nu}^{ab}(k) &= -i g_{\mu\nu} (k^2 - m_a^2) \delta_{ab} - i \left(g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) \hat{\Sigma}_T^{ab}(k^2) - i \frac{k_\mu k_\nu}{k^2} \hat{\Sigma}_L^{ab}(k^2)
\end{aligned} \tag{3.3}$$

Here, $a, b = \gamma, Z$. From the above equations, we obtain [23]

$$\begin{aligned}
\delta m_W^2 &= \tilde{R}e \Sigma_T^{WW}(m_W^2), & \delta m_Z^2 &= \tilde{R}e \Sigma_T^{ZZ}(m_Z^2) \\
\delta Z_W &= -\tilde{R}e \frac{\partial \Sigma_T^{WW}(k^2)}{\partial k^2} \big|_{k^2=m_W^2} \\
\delta Z_e &= -\frac{1}{2} \delta Z_{\gamma\gamma} - \frac{s_W}{c_W} \frac{1}{2} \delta Z_{Z\gamma} = \frac{1}{2} \frac{\partial \Sigma_T^{\gamma\gamma}(k^2)}{\partial k^2} \big|_{k^2=0} - \frac{s_W}{c_W} \frac{\Sigma_T^{\gamma\gamma}(0)}{m_Z^2},
\end{aligned} \tag{3.4}$$

in which Σ 's are the corresponding unrenormalized self energies.

By using the equation $\cos^2 \theta_W = m_W^2/m_Z^2$, we find

$$\delta \cos \theta_W = \frac{1}{2} \cos \theta_W \left(\frac{\delta m_W^2}{m_W^2} - \frac{\delta m_Z^2}{m_Z^2} \right). \tag{3.5}$$

(b) The renormalized one-particle-irreducible two-point Green functions of fermions (here they are charginos and neutralinos) can be decomposed as

$$i\Gamma_{ij}^f(p) = i\delta_{ij}(\not{p} - m_{f_i}) + i \left[\not{p}P_L\hat{\Sigma}_{ij}^L + \not{p}P_R\hat{\Sigma}_{ij}^R + P_L\hat{\Sigma}_{ij}^{SL} + P_R\hat{\Sigma}_{ij}^{SR} \right]. \quad (3.6)$$

where $\hat{\Sigma}_{ij}^{L,R,SL,SR}$ are the renormalized self-energy matrices of fermions. By imposing the on-mass-shell renormalization conditions for fermions [23]

$$\begin{aligned} \lim_{p^2 \rightarrow m_{f_i}^2} \frac{1}{i(\not{p} - m_{f_i})} \tilde{R}e i \hat{\Gamma}_{ii}^f(p) u_i(p) &= u_i(p) & \tilde{R}e i \hat{\Gamma}_{ij}^f(p) u_j(p) \big|_{p^2=m_{f_j}^2} &= 0 \\ \lim_{p^2 \rightarrow m_{f_i}^2} \bar{u}_i(p) \tilde{R}e i \hat{\Gamma}_{ii}^f(p) \frac{1}{i(\not{p} - m_{f_i})} &= \bar{u}_i(p) & \tilde{R}e \bar{u}_i(p) i \hat{\Gamma}_{ij}^f(p) \big|_{p^2=m_{f_i}^2} &= 0 \end{aligned} \quad (3.7)$$

we can get the renormalization constants of fermions [23] [25] [24]:

$$\begin{aligned} \delta m_i &= \frac{1}{2} Re \left[m_i (\Sigma_{ii}^L(m_i^2) + \Sigma_{ii}^R(m_i^2)) + \Sigma_{ii}^{SL}(m_i^2) + \Sigma_{ii}^{SR}(m_i^2) \right], \\ \delta Z_{ii}^L &= -Re \left[\Sigma_{ii}^L(m_i^2) + m_i^2 (\Sigma_{ii}^{L'}(m_i^2) + \Sigma_{ii}^{R'}(m_i^2)) + m_i (\Sigma_{ii}^{SL'}(m_i^2) + \Sigma_{ii}^{SR'}(m_i^2)) \right], \\ \delta Z_{ii}^R &= -Re \left[\Sigma_{ii}^R(m_i^2) + m_i^2 (\Sigma_{ii}^{L'}(m_i^2) + \Sigma_{ii}^{R'}(m_i^2)) + m_i (\Sigma_{ii}^{SL'}(m_i^2) + \Sigma_{ii}^{SR'}(m_i^2)) \right], \\ \delta Z_{ij}^L &= \frac{2}{m_i^2 - m_j^2} Re \left[m_j^2 \Sigma_{ij}^L(m_j^2) + m_i m_j \Sigma_{ij}^R(m_j^2) + m_i \Sigma_{ij}^{SL}(m_j^2) + m_j \Sigma_{ij}^{SR}(m_j^2) \right], \quad \text{for } i \neq j \\ \delta Z_{ij}^R &= \frac{2}{m_i^2 - m_j^2} Re \left[m_i m_j \Sigma_{ij}^L(m_j^2) + m_j^2 \Sigma_{ij}^R(m_j^2) + m_j \Sigma_{ij}^{SL}(m_j^2) + m_i \Sigma_{ij}^{SR}(m_j^2) \right]. \quad \text{for } i \neq j \end{aligned} \quad (3.8)$$

For radiative corrections to chargino and neutralino masses, we can refer to Ref.[26].

(c) $\delta U, \delta V$ and δN are fixed by demanding that the wave function corrections are symmetric [25] [24]:

$$\begin{aligned} \delta U &= \frac{1}{4} \left(\delta Z_+^R - \delta Z_+^{R\dagger} \right) U \\ \delta V &= \frac{1}{4} \left(\delta Z_+^L - \delta Z_+^{L\dagger} \right) V \end{aligned}$$

$$\delta N = \frac{1}{4} \left(\delta Z_0^L - \delta Z_0^{L\dagger} \right) N \quad (3.9)$$

The chargino and neutralino self-energies can be found in Ref.[24].

By substituting Eq.(3.1) into the Lagrangian (2.9), we can obtain the counter-term of vertex $\bar{\tilde{\chi}}_i^0 \tilde{\chi}_j^+ W^-$:

$$\begin{aligned} \delta \mathcal{L}_{\tilde{\chi}^+ \tilde{\chi}^0 W} = & g W_\mu^- \bar{\tilde{\chi}}_i^0 \gamma^\mu \left\{ P_L \left[\delta O_{ij}^L + O_{ij}^L \left(\frac{\delta g}{g} + \frac{1}{2} \delta Z_W \right) + \frac{1}{2} \sum_{k=1}^2 O_{ik}^L \delta Z_{+,kj}^L \right. \right. \\ & + \frac{1}{2} \sum_{k=1}^4 O_{kj}^L \delta Z_{0,ki}^{L*} \left. \right] + P_R \left[\delta O_{ij}^R + O_{ij}^R \left(\frac{\delta g}{g} + \frac{1}{2} \delta Z_W \right) + \frac{1}{2} \sum_{k=1}^2 O_{ik}^R \delta Z_{+,kj}^R \right. \\ & \left. \left. + \frac{1}{2} \sum_{k=1}^4 O_{kj}^R \delta Z_{0,ki}^{R*} \right] \right\} \tilde{\chi}_j^+ + h.c. \end{aligned} \quad (3.10)$$

where

$$\begin{aligned} \delta O_{ij}^L &= -\frac{1}{\sqrt{2}} (\delta N_{i4} V_{j2}^* + N_{i4} \delta V_{j2}^*) + (\delta N_{i2} V_{j1}^* + N_{i2} \delta V_{j1}^*) \\ \delta O_{ij}^R &= \frac{1}{\sqrt{2}} (\delta N_{i3}^* U_{j2} + N_{i3}^* \delta U_{j2}) + (\delta N_{i2}^* U_{j1} + N_{i2}^* \delta U_{j1}) \end{aligned} \quad (3.11)$$

For convenience, we define

$$\begin{aligned} -i\delta \Lambda_{ji}^L &= O_{ij}^L \left(\frac{\delta g}{g} + \frac{1}{2} \delta Z_W \right) + \frac{1}{2} \sum_{k=1}^2 O_{ik}^L \delta Z_{+,kj}^L + \frac{1}{2} \sum_{k=1}^4 O_{kj}^L \delta Z_{0,ki}^{L*} + \delta O_{ij}^L \\ -i\delta \Lambda_{ji}^R &= O_{ij}^R \left(\frac{\delta g}{g} + \frac{1}{2} \delta Z_W \right) + \frac{1}{2} \sum_{k=1}^2 O_{ik}^R \delta Z_{+,kj}^R + \frac{1}{2} \sum_{k=1}^4 O_{kj}^R \delta Z_{0,ki}^{R*} + \delta O_{ij}^R \end{aligned} \quad (3.12)$$

Then the contribution from the counter-term to the amplitude of the decay $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$ can be written as

$$\delta \mathcal{M}_c = \bar{u}_{\tilde{\chi}_1^0}(k_1) \gamma^\mu (\delta \Lambda_{11}^L P_L + \delta \Lambda_{11}^R P_R) u_{\tilde{\chi}_1^+}(p_1) \epsilon_\mu(k_2) \quad (3.13)$$

Up to the one-loop level, the total amplitude of this process is

$$\mathcal{M}_{tot} = \mathcal{M}_{tree} + \delta\mathcal{M}_v + \delta\mathcal{M}_c \quad (3.14)$$

where $\delta\mathcal{M}_v$ represents the vertex correction. For regularization of the ultraviolet divergences in the virtual loop corrections, we adopt the dimensional reduction scheme (DR) [27], which is commonly used in the calculations of the electroweak corrections in framework of the MSSM as it preserves supersymmetry at least at one-loop level. As we expected, the divergence in $\delta\mathcal{M}_v$ will be cancelled by that in the counter-term $\delta\mathcal{M}_c$ exactly. We can check this property of renormalization both analytically and numerically.

The calculations of the vertex correction $\delta\mathcal{M}_v$ contributed by the one-loop diagrams in Fig.1(c) tell us that

$$\delta\mathcal{M}_v = \bar{u}_{\tilde{\chi}_1^0}(k_1)[\gamma^\mu(\Lambda_{11}^L P_L + \Lambda_{11}^R P_R) + p_1^\mu(\Pi_{11}^L P_L + \Pi_{11}^R P_R)]u_{\tilde{\chi}_1^+}(p_1)\epsilon_\mu(k_2) \quad (3.15)$$

where Λ_{11}^L , Λ_{11}^R , Π_{11}^L and Π_{11}^R are the form factors whose expressions are listed in the Appendix. To describe the magnitude of the supersymmetric electroweak corrections to the partial width of the decay $\tilde{\chi}_1^+ \rightarrow W^+\tilde{\chi}_1^0$, we introduce a quantity named relative correction δ defined as

$$\delta = \frac{\Gamma_{tot}(\tilde{\chi}_1^+ \rightarrow W^+\tilde{\chi}_1^0) - \Gamma_{tree}(\tilde{\chi}_1^+ \rightarrow W^+\tilde{\chi}_1^0)}{\Gamma_{tree}(\tilde{\chi}_1^+ \rightarrow W^+\tilde{\chi}_1^0)} \quad (3.16)$$

It will be used in numerical calculations in the next section.

IV Numerical results and conclusion

In the following we present some numerical results for the radiative corrections to chargino decay into the lightest neutralino and W boson. In our numerical calculations the SM input

parameters are chosen to be $m_t = 174.3$ GeV, $m_b = 4.3$ GeV, $m_Z = 91.188$ GeV, $m_W = 80.41$ GeV and $\alpha_{EW} = 1/128$ [28]. For the SUSY sector, we choose the physical observables to be the input parameters rather than the original parameters in the Lagrangian (e.g. the soft-SUSY-breaking parameters $M, M', M_{\tilde{Q}}^2, \dots$). In this paper we use the following set of independent parameters:

$$(\tan \beta, m_{\tilde{\chi}_1^+}, m_{\tilde{\chi}_2^+}, m_{\tilde{\chi}_1^0}, m_{\tilde{t}_1}, \theta_t, m_{\tilde{b}_1}, m_{\tilde{b}_2}, \theta_b) \quad (4.1)$$

The parameters $\tan \beta, m_{\tilde{\chi}_1^+}, m_{\tilde{\chi}_2^+}$ and $m_{\tilde{\chi}_1^0}$ among these nine input parameters are used to determine the chargino and neutralino sectors and the rest five parameters to determine the squark sectors. The parameters μ and M in the chargino matrix X (see eq.(2.2)) can be extracted from the input chargino masses $m_{\tilde{\chi}_{1,2}^+}$, but their values are not uniquely determined. For the fixed chargino masses, we can obtain four sets of results:

case 1: $-\mu > M > 0$, case2: $M > -\mu > 0$, case 3: $M > \mu > 0$ and case 4: $\mu > M > 0$.

The only differences among the four cases present in their U and V matrices. All the four cases are considered in this paper for completeness. The value of $m_{\tilde{t}_2}$ is determined by $SU(2)$ gauge invariance. For all the numerical calculations in this paper, we take $m_{\tilde{\chi}_2^+} = 550$ GeV, $m_{\tilde{b}_1} = 400$ GeV, $m_{\tilde{b}_2} = 450$ GeV and $m_{\tilde{t}_1} = 140$ GeV.

In Fig.2 we present the relative correction δ as a function of $\tan \beta$ for $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$, assuming $m_{\tilde{\chi}_1^+} = 190$ GeV, $m_{\tilde{\chi}_1^0} = 90$ GeV and $(\theta_b, \theta_t) = (-0.3, 0.78), (-0.2, 0.4), (0.2, -0.4), (0.35, -0.4)$ for case 1, 2, 3 and 4 respectively. As shown in this figure, the SUSY-EW corrections are important and especially sensitive to the value of $\tan \beta$ for case 1 and 4. For case 1 the relative correction δ is positive when $\tan \beta \gtrsim 4$. It varies from -5% to 11.9% as the increment of $\tan \beta$ from 2 to 9 and decreases slowly as $\tan \beta$ grows in the

region where $\tan\beta > 9$. The peak at $\tan\beta \sim 9$ on this curve is due to the fact that the tree-level partial width of this decay Γ_{tree} has a minimum value around this position. For case 4 the relative correction is negative, large ($|\delta| > 11\%$) and increases to -16.6% as $\tan\beta \sim 36$. For case 2 and 3, the relative corrections are insensitive to $\tan\beta$. They are all negative and vary from -8.4% to -10% and -9.8% to -11.1% , respectively.

Fig.3(a) and (b) give the relative corrections as the functions of $m_{\tilde{\chi}_1^+}$ in the case of $m_{\tilde{\chi}_1^0} = 60$ GeV and 90 GeV, respectively, assuming $\tan\beta = 7$. The values of (θ_b, θ_t) are the same as in Fig.2. Both in the case of $m_{\tilde{\chi}_1^0} = 60$ GeV and 90 GeV, the relative corrections are insensitive with $m_{\tilde{\chi}_1^+}$ for case 2 and 3. The variations of them are all less than 1%. Furthermore, the curves for case 2, 3 and 4 show that the relative corrections in Fig.3(a) are almost the same as in Fig.3(b) for a fixed $m_{\tilde{\chi}_1^+}$. This conclusion can be obtained from Fig.4 more evidently. The corrections are about -9% and -10% for case 2 and 3 respectively and vary from -13.7% to about -10% for case 4. For case 1, the relative corrections range between 9% (11.3%) and 3.7% (1.7%) with $m_{\tilde{\chi}_1^+}$ varying in the mass region 150 (180) GeV to 270 (280) GeV when $m_{\tilde{\chi}_1^0} = 60$ (90) GeV.

In Fig.4 we show graphically the dependence between the relative correction and $m_{\tilde{\chi}_1^0}$. Here we take $\tan\beta = 7$, $m_{\tilde{\chi}_1^+} = 190$ GeV. θ_b and θ_t are set to be the same as in figure 2. It is clear that the relative corrections are rather stable as $m_{\tilde{\chi}_1^0}$ running from 60 GeV to 100 GeV for case 2, 3, 4 and are also not very insensitive to $m_{\tilde{\chi}_1^0}$ for case 1. It can reach about 11.7% and -13.4% for case 1 and 4, respectively.

The relative corrections as the functions of the squark mixing angles θ_b and θ_t are displayed in Fig.5 and Fig.6, respectively. The insensitivity of the relative correction δ to

θ_b is due to the fact that the sbottoms $\tilde{b}_{1,2}$ ($m_{\tilde{b}_{1,2}} = 400, 450$ GeV) are much heavier than the lighter stop $\tilde{t}_1$ ($m_{\tilde{t}_1} = 140$ GeV). Therefore, the sbottom quark are almost decoupled and the variation of θ_b does not affect the relative correction too much. In contrast to the sbottom mixing angle θ_b , the stop mixing angle θ_t plays a crucial role, as seen in Fig.6. The SUSY-EW corrections decrease or increase the partial decay width of $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$ significantly depending on θ_t .

In summary, we have computed the supersymmetric electroweak corrections to the partial width of the decay $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$ in the MSSM. Only the third generation of quark and squark are considered because of the Yukawa-like couplings of the relevant vertices. These corrections are not very sensitive to the mass of the lightest neutralino and the sbottom mixing angle, but depend strongly on the parameters $\tan\beta$ and θ_t . The magnitude of the corrections can exceed 10% in some parameter space, therefore they should be taken into account in any reliable analysis.

Acknowledgements: This work was supported in part by the National Natural Science Foundation of China(project Nos: 19875049, 10005009), the Doctoral Program Foundation of the Education Ministry of China, and a grant from the Ministry of Science and Technology of China.

V Appendix

In this appendix, we list the form factors of the decay $\tilde{\chi}_i^+ \rightarrow W^+ \tilde{\chi}_j^0$ ($i=1,2, j=1,2,3,4$). The form factors $\Lambda_{ij}^{L,R}$ and $\Pi_{ij}^{L,R}$ can be divided into four parts respectively

$$\Lambda_{ij}^{L,R} = \Lambda_{ij}^{(1)L,R} + \Lambda_{ij}^{(2)L,R} + \Lambda_{ij}^{(3)L,R} + \Lambda_{ij}^{(4)L,R}$$

$$\Pi_{ij}^{L,R} = \Pi_{ij}^{(1)L,R} + \Pi_{ij}^{(2)L,R} + \Pi_{ij}^{(3)L,R} + \Pi_{ij}^{(4)L,R} \quad (5.1)$$

where $\Lambda_{ij}^{(a)L,R}$ and $\Pi_{ij}^{(a)L,R}$ ($a=1,2,3,4$) are the form factors contributed by the four one-loop diagrams in Fig.1($c-1 \sim c-4$), respectively. In the calculation of these form factors, the following vertices (or their conjugate vertices) will be used:

$$\begin{aligned} \bar{t} - \tilde{\chi}_i^+ - \tilde{b}_j & : & V_{t\tilde{\chi}_i^+\tilde{b}_j}^L P_L + V_{t\tilde{\chi}_i^+\tilde{b}_j}^R P_R \\ \bar{b} - \tilde{\chi}_i^+ - \tilde{t}_j & : & (V_{b\tilde{\chi}_i^+\tilde{t}_j}^L P_L + V_{b\tilde{\chi}_i^+\tilde{t}_j}^R P_R) C \\ \bar{t} - \tilde{\chi}_i^0 - \tilde{t}_j & : & V_{t\tilde{\chi}_i^0\tilde{t}_j}^L P_L + V_{t\tilde{\chi}_i^0\tilde{t}_j}^R P_R \\ \bar{b} - \tilde{\chi}_i^0 - \tilde{b}_j & : & V_{b\tilde{\chi}_i^0\tilde{b}_j}^L P_L + V_{b\tilde{\chi}_i^0\tilde{b}_j}^R P_R \\ \tilde{b}_i^* - \tilde{t}_j - W^- & : & V_{\tilde{b}_i\tilde{t}_jW} (p + p')^\mu \end{aligned} \quad (5.2)$$

where p and p' are the incoming momentum of $\tilde{t}_j$ and outgoing momentum of $\tilde{b}_i$ respectively. The explicit expressions of these vertices can be found in Ref.[3]. and [20].

Now we can write down the form factors of the decay $\tilde{\chi}_i^+(p_1) \rightarrow \tilde{\chi}_j^0(k_1) + W^+(k_2)$:

$$\begin{aligned} \Lambda_{ij}^{(1)L} &= \frac{3}{8\pi^2} \sum_{l,k=1}^2 V_{b\tilde{\chi}_i^+\tilde{t}_l}^{R*} V_{b\tilde{\chi}_j^0\tilde{b}_k}^R V_{\tilde{b}_k\tilde{t}_lW} C_{24}^{(1)} \\ \Pi_{ij}^{(1)L} &= -\frac{3}{8\pi^2} \sum_{l,k=1}^2 V_{\tilde{b}_k\tilde{t}_lW} [(C_{11}^{(1)} - C_{12}^{(1)} + C_{21}^{(1)} - C_{23}^{(1)}) V_{b\tilde{\chi}_i^+\tilde{t}_l}^{R*} V_{b\tilde{\chi}_j^0\tilde{b}_k}^R m_{\tilde{\chi}_j^0} \\ &\quad + (C_{11}^{(1)} - C_{12}^{(1)}) V_{b\tilde{\chi}_i^+\tilde{t}_l}^{R*} V_{b\tilde{\chi}_j^0\tilde{b}_k}^L m_b + (C_{22}^{(1)} - C_{23}^{(1)}) V_{b\tilde{\chi}_i^+\tilde{t}_l}^{L*} V_{b\tilde{\chi}_j^0\tilde{b}_k}^L m_{\tilde{\chi}_i^+}] \\ \Lambda_{ij}^{(1)R} &= \Lambda_{ij}^{(1)L} (L \leftrightarrow R) \\ \Pi_{ij}^{(1)R} &= \Pi_{ij}^{(1)L} (L \leftrightarrow R) \\ \Lambda_{ij}^{(2)L} &= -\frac{3}{8\pi^2} \sum_{l,k=1}^2 V_{t\tilde{\chi}_i^+\tilde{b}_k}^L V_{t\tilde{\chi}_j^0\tilde{t}_l}^{L*} V_{\tilde{b}_k\tilde{t}_lW} C_{24}^{(2)} \\ \Pi_{ij}^{(2)L} &= \frac{3}{8\pi^2} \sum_{l,k=1}^2 V_{\tilde{b}_k\tilde{t}_lW} [(C_{11}^{(2)} - C_{12}^{(2)} + C_{21}^{(2)} - C_{23}^{(2)}) V_{t\tilde{\chi}_i^+\tilde{b}_k}^L V_{t\tilde{\chi}_j^0\tilde{t}_l}^{L*} m_{\tilde{\chi}_j^0} \end{aligned}$$

$$\begin{aligned}
& +(C_{11}^{(2)} - C_{12}^{(2)})V_{t\tilde{\chi}_i^+\tilde{b}_k}^L V_{t\tilde{\chi}_j^0\tilde{t}_l}^{R*} m_t + (C_{22}^{(2)} - C_{23}^{(2)})V_{t\tilde{\chi}_i^+\tilde{b}_k}^R V_{t\tilde{\chi}_j^0\tilde{t}_l}^{R*} m_{\tilde{\chi}_i^+}] \\
\Lambda_{ij}^{(2)R} &= \Lambda_{ij}^{(2)L}(L \leftrightarrow R) \\
\Pi_{ij}^{(2)R} &= \Pi_{ij}^{(2)L}(L \leftrightarrow R) \\
\Lambda_{ij}^{(3)L} &= -i\frac{3g}{16\sqrt{2}\pi^2} \sum_{k=1}^2 \{V_{t\tilde{\chi}_i^+\tilde{b}_k}^L m_t [C_0^{(3)} m_b V_{b\tilde{\chi}_j^0\tilde{b}_k}^{L*} - C_{11}^{(3)} m_{\tilde{\chi}_j^0} V_{b\tilde{\chi}_j^0\tilde{b}_k}^{R*}] \\
& + V_{t\tilde{\chi}_i^+\tilde{b}_k}^R m_{\tilde{\chi}_i^+} [(C_0^{(3)} + C_{12}^{(3)}) m_b V_{b\tilde{\chi}_j^0\tilde{b}_k}^{L*} - (C_{11}^{(3)} - C_{12}^{(3)}) m_{\tilde{\chi}_j^0} V_{b\tilde{\chi}_j^0\tilde{b}_k}^{R*}] \} \\
\Lambda_{ij}^{(3)R} &= i\frac{3g}{16\sqrt{2}\pi^2} \sum_{k=1}^2 \{C_{12}^{(3)} m_t m_{\tilde{\chi}_i^+} V_{t\tilde{\chi}_i^+\tilde{b}_k}^L V_{b\tilde{\chi}_j^0\tilde{b}_k}^{R*} - (C_0^{(3)} + C_{11}^{(3)}) m_b m_{\tilde{\chi}_j^0} V_{t\tilde{\chi}_i^+\tilde{b}_k}^R V_{b\tilde{\chi}_j^0\tilde{b}_k}^{L*} \\
& + [\frac{1}{2} - 2C_{24}^{(3)} + (C_{12}^{(3)} + C_{23}^{(3)}) m_W^2 + (C_{22}^{(3)} - C_{23}^{(3)}) m_{\tilde{\chi}_i^+}^2 \\
& + (C_{11}^{(3)} - C_{12}^{(3)} + C_{21}^{(3)} - C_{23}^{(3)}) m_{\tilde{\chi}_j^0}^2] V_{t\tilde{\chi}_i^+\tilde{b}_k}^R V_{b\tilde{\chi}_j^0\tilde{b}_k}^{R*} \} \\
\Pi_{ij}^{(3)L} &= -i\frac{3g}{8\sqrt{2}\pi^2} \sum_{k=1}^2 [C_{12}^{(3)} m_t V_{t\tilde{\chi}_i^+\tilde{b}_k}^L + (C_{22}^{(3)} - C_{23}^{(3)}) m_{\tilde{\chi}_i^+} V_{t\tilde{\chi}_i^+\tilde{b}_k}^R] V_{b\tilde{\chi}_j^0\tilde{b}_k}^{R*} \\
\Pi_{ij}^{(3)R} &= i\frac{3g}{8\sqrt{2}\pi^2} \sum_{k=1}^2 V_{t\tilde{\chi}_i^+\tilde{b}_k}^R [(C_0^{(3)} + C_{11}^{(3)}) m_b V_{b\tilde{\chi}_j^0\tilde{b}_k}^{L*} - (C_{11}^{(3)} - C_{12}^{(3)} + C_{21}^{(3)} - C_{23}^{(3)}) m_{\tilde{\chi}_j^0} V_{b\tilde{\chi}_j^0\tilde{b}_k}^{R*}] \\
\Lambda_{ij}^{(4)L} &= -i\frac{3g}{16\sqrt{2}\pi^2} \sum_{k=1}^2 \{C_{12}^{(4)} m_b m_{\tilde{\chi}_i^+} V_{b\tilde{\chi}_i^+\tilde{t}_k}^{L*} V_{t\tilde{\chi}_j^0\tilde{t}_k}^R - (C_0^{(4)} + C_{11}^{(4)}) m_t m_{\tilde{\chi}_j^0} V_{b\tilde{\chi}_i^+\tilde{t}_k}^{R*} V_{t\tilde{\chi}_j^0\tilde{t}_k}^L \\
& + [\frac{1}{2} - 2C_{24}^{(4)} + (C_{12}^{(4)} + C_{23}^{(4)}) m_W^2 + (C_{22}^{(4)} - C_{23}^{(4)}) m_{\tilde{\chi}_i^+}^2 \\
& + (C_{11}^{(4)} - C_{12}^{(4)} + C_{21}^{(4)} - C_{23}^{(4)}) m_{\tilde{\chi}_j^0}^2] V_{b\tilde{\chi}_i^+\tilde{t}_k}^{R*} V_{t\tilde{\chi}_j^0\tilde{t}_k}^R \} \\
\Lambda_{ij}^{(4)R} &= i\frac{3g}{16\sqrt{2}\pi^2} \sum_{k=1}^2 \{V_{b\tilde{\chi}_i^+\tilde{t}_k}^{L*} m_b [C_0^{(4)} m_t V_{t\tilde{\chi}_j^0\tilde{t}_k}^L - C_{11}^{(4)} m_{\tilde{\chi}_j^0} V_{t\tilde{\chi}_j^0\tilde{t}_k}^R] \\
& + V_{b\tilde{\chi}_i^+\tilde{t}_k}^{R*} m_{\tilde{\chi}_i^+} [(C_0^{(4)} + C_{12}^{(4)}) m_t V_{t\tilde{\chi}_j^0\tilde{t}_k}^L - (C_{11}^{(4)} - C_{12}^{(4)}) m_{\tilde{\chi}_j^0} V_{t\tilde{\chi}_j^0\tilde{t}_k}^R] \} \\
\Pi_{ij}^{(4)L} &= -i\frac{3g}{8\sqrt{2}\pi^2} \sum_{k=1}^2 V_{b\tilde{\chi}_i^+\tilde{t}_k}^{R*} [(C_0^{(4)} + C_{11}^{(4)}) m_t V_{t\tilde{\chi}_j^0\tilde{t}_k}^L - (C_{11}^{(4)} - C_{12}^{(4)} + C_{21}^{(4)} - C_{23}^{(4)}) m_{\tilde{\chi}_j^0} V_{t\tilde{\chi}_j^0\tilde{t}_k}^R] \\
\Pi_{ij}^{(4)R} &= i\frac{3g}{8\sqrt{2}\pi^2} \sum_{k=1}^2 [C_{12}^{(4)} m_b V_{b\tilde{\chi}_i^+\tilde{t}_k}^{L*} + (C_{22}^{(4)} - C_{23}^{(4)}) m_{\tilde{\chi}_i^+} V_{b\tilde{\chi}_i^+\tilde{t}_k}^{R*}] V_{t\tilde{\chi}_j^0\tilde{t}_k}^R \tag{5.3}
\end{aligned}$$

where

$$\begin{aligned}
C_{11,12,21,22,23,24}^{(1)} &= C_{11,12,21,22,23,24}(k_1, -p_1, m_{\tilde{b}_k}, m_b, m_{\tilde{t}_l}) \\
C_{11,12,21,22,23,24}^{(2)} &= C_{11,12,21,22,23,24}(k_1, -p_1, m_{\tilde{t}_l}, m_t, m_{\tilde{b}_k}) \\
C_{0,11,12,21,22,23,24}^{(3)} &= C_{0,11,12,21,22,23,24}(k_1, -p_1, m_b, m_{\tilde{b}_k}, m_t) \\
C_{0,11,12,21,22,23,24}^{(4)} &= C_{0,11,12,21,22,23,24}(k_1, -p_1, m_t, m_{\tilde{t}_k}, m_b) \tag{5.4}
\end{aligned}$$

The definitions and numerical calculation formula of the one-point, two-point and three-point Passarino-Veltman integrals are adopted from Ref.[29] and Ref.[30], respectively.

References

- [1] S. Weinberg, Phys. Rev. Lett. **19** (1967) 1264; S. Glashow, Nucl. Phys. **22** (1961) 579; A. Salam, in *Elementary Particle Theory*, ed. N. Svartholm, (1968) p367.
- [2] H. P. Nilles Phys. Rep. **110** (1984) 1; R. Barbieri, Riv. Nuov. Cim. **11** 1 (1988).
- [3] H. E. Haber, G. L. Kane, Phys. Rep. **117** (1985) 75.
- [4] A. B. Lahanas, D. V. Nanopoulos, Phys. Rep. **145** (1987) 1.
- [5] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429** (1998) 263, hep-ph/9803315; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett.**B436** (1998) 257, hep-ph/9804398; N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Rev. **D59** (1999) 086004, hep-ph/9807344.
- [6] OPAL Collab., K. Ackerstaff, *et al.*, Eur. Phys. J. **C6** (1999) 225; G. Abbiendi, *et al.*, Phys. Lett. **B456** (1999) 95; L3 Collab., Phys. Lett. **B471** (1999) 308.

- [7] D0 Collab., B. Abbott, *et al.*, Phys. Rev. **D60** (1999) 031101; Phys. Rev. Lett. **83** (1999) 4937; V. M. Abazov, *et al.*, hep-ex/0108018; CDF Collab., hep-ex/0106001; hep-ex/0106061; Phys. Rev. **D63** (2001) 091101; Phys. Rev. Lett. **84** (2000) 5273.
- [8] M. Spiropulu, for the D0 Collaboration, the CDF Collaboration, EFI-01-23, FNALPUB-Conf-01-110-E, hep-ex/0106045.
- [9] OPAL Collab., K. Ackerstaff, *et al.*, Eur. Phys. J. **C2** (1998) 213; G. Abbiendi, *et al.*, Eur. Phys. J. **C8** (1999) 255; DELPHI Collab., P. Abreu, *et al.*, Phys. Lett. **B466** (1999) 75; Eur. Phys. J. **C14** (2000) 187; Erratum-ibid. **C16** (2000) 707; L3 Collab., Phys. Lett. **B482** (2000) 31; Phys. Lett. **B472** (2000) 420.
- [10] CDF Collab., F. Abe, *et al.*, Phys. Rev. Lett. **76** (1996) 4307; Phys. Rev. Lett. **80** (1998) 5275.
- [11] Masayuki Koike, Toshihiko Nonaka, Tadashi Kon, Phys. Lett. **B357** (1995) 232, hep-ph/9504309.
- [12] M. L. Zhou, W. G. Ma, L. Han, Y. Jiang, H. Zhou, J. Phys. **G25** (1999) 1641, hep-ph/9903376.
- [13] S. Hesselbach, H. Fraas, Phys. Rev. **D55** (1997) 1343-1352, hep-ph/9604439.
- [14] Marco A. Diaz, Steve F. King, Douglas A. Ross, hep-ph/9801373, hep-ph/0012340; S. Y. Choi, A. Djouadi, H. Dreiner, J. Kalinowski, P. M. Zerwas, Eur. Phys. J. **C7** (1999) 123-134, hep-ph/9806279; A. Bartl, H. Eberl, H. Fraas, S. Kraml, W.

- Majerotto, G. Moortgat-Pick, W. Porod, hep-ph/0010018; T. Blank, W. Hollik, hep-ph/0011092.
- [15] M. Czakon, J. Gluza, hep-ph/0003228.
- [16] Z. Z. Aydin, O. Yilmaz, Europhys. Lett. **50** (2000) 22-27 hep-ph/0002206.
- [17] W. G. Ma, F. Du, M. L. Zhou, Y. Jiang, L. Han, Phys. Rev. **D60** (1999) 115009, hep-ph/9907368; L. H. Wan, W. G. Ma, X. Yin, Y. Jiang, L. Han, J. Phys. **G27** (2001) 203 hep-ph/0012078; Y. Jang, W. G. Ma, L. Han, Z. H. Yu, H. Pietschmann, Phys. Rev. **D62** (2000) 035006.
- [18] A. Djouadi, Y. Mambrini, M. Muhlleitner, Eur. Phys. J. **C20** (2001) 563.
- [19] J. L. Feng, M. J. Strassler, Phys. Rev. **D55** (1997) 1326.
- [20] J. F. Gunion, H. E. Haber, Nucl. Phys. **B272** (1986) 1.
- [21] S. Kiyoura, M. Nojiri, D. M. Pierce, Y. Yamada, Phys. Rev. **D58** (1998) 075002.
- [22] M. A. Diaz, D. A. Ross, JHEP **0106** (2001) 001.
- [23] A. Denner, Fortschr. Phys. **41** (1993) 307.
- [24] H. Eberl, M. Kincel, W. Majerotto, Y. Yamada, hep-ph/0104109.
- [25] A. Denner, T. Sack, Nucl. Phys. **B347** 203 (1990); Y. Yamada, Phys. Rev. **D64** (2001) 036008, hep-ph/0103046.
- [26] D. Pierce, A. Papadopoulos, Phys. Rev. **D50** (1994) 565.

- [27] D. M. Copper, D. R. T. Jones, P. van Nieuwennuizen, Nucl. Phys. **B167** (1980) 479;
W. Siegel, Phys. Lett. **B84** (1979) 193.
- [28] Particle Data Group, Eur. Phys. J. **C15** (2000) 1.
- [29] Bernd A. Kniehl, Phys. Rep. **240** (1994) 211.
- [30] G. Passarino and M. Veltman, Nucl. Phys. **B160**, (1979) 151.

Figure Captions

Fig.1 The relevant Feynman diagrams to the decay $\tilde{\chi}_1^+ \rightarrow W^+ \tilde{\chi}_1^0$: (a) tree-level diagram; (b) counter-term for vertex; (c-1)-(c-4) one-loop vertex corrections; (d), (e) and (f) self-energies of W boson, chargino and neutralino respectively. In diagrams (a), (b) and (c), $i = j = 1$. In diagrams (c), (d), (e) and (f) the subscript k and l can take from 1 to 2.

Fig.2 The relative correction as a function of $\tan \beta$.

Fig.3 The relative correction as a function of $m_{\tilde{\chi}_1^+}$.

Fig.4 The relative correction as a function of $m_{\tilde{\chi}_1^0}$.

Fig.5 The relative correction as a function of θ_b .

Fig.6 The relative correction as a function of θ_t .




Fig.1

Fig.2




Fig.3 (a)




Fig.3 (b)




Fig.4




Fig.5




Fig.6

